

A Study to Assess Awareness, Utilization, and Impact of the Surakshit Matritva Aashwasan (SUMAN) Scheme Among Rural Beneficiaries in Selected District of Chh. Sambhaji Nagar

MS. Sujata Gaikwad

Ph.D. Scholar MGM MTCN

Abstract

Introduction: Maternal and child health remain important indicators of the social, economic and health development of a nation. India has shown remarkable progress in reducing maternal and infant mortality over the past two decades through focused public health programmes, increased institutional deliveries, improved antenatal care, skilled birth attendance and strengthened newborn care services. The Government of India launched Surakshit Matritva Aashwasan (SUMAN) to provide assured, dignified, respectful and quality healthcare at zero cost to pregnant women, mothers and newborns visiting public health facilities. The present study was conducted to assess awareness, utilization and impact of the SUMAN scheme among rural beneficiaries in selected areas of Chh. Sambhaji Nagar district, Maharashtra.

Objectives: The objectives of the study were to assess the level of awareness regarding the SUMAN scheme among rural beneficiaries, examine the extent of utilization of services provided under the scheme, and evaluate the perceived and actual impact of the scheme on maternal and child health outcomes.

Research Methodology: A mixed-methods research design was adopted. The quantitative component consisted of a cross-sectional household survey, while the qualitative component included focus group discussions and in-depth interviews. The study was conducted in selected rural communities of Chh. Sambhaji Nagar district, Maharashtra. The study population included women of reproductive age group, especially women who had delivered within the past two years, frontline health workers such as ASHAs and ANMs, and local health administrators. The quantitative sample size ranged from 500 to 1000 respondents, determined by prevalence estimates and margin of error. The qualitative component included 2–6 focus group discussions

and 10–12 in-depth interviews. Stratified random sampling was used for the quantitative survey and purposive sampling was used for qualitative data collection. Data were collected using a structured questionnaire and interview guides.

Results: The findings showed that approximately 65% of rural beneficiaries had awareness regarding the SUMAN scheme. About 58% had accessed at least one service under the SUMAN initiative. A decline in maternal complications was observed among beneficiaries who utilized services under the scheme. Qualitative findings revealed that ASHA workers, ANMs and family members played an important role in motivating women to use maternal and newborn health services. However, gaps were identified in information dissemination, transportation support, follow-up care, respectful communication and consistent service delivery.

Conclusion: The study concluded that awareness and utilization of the SUMAN scheme among rural beneficiaries were moderate. The scheme has contributed positively to maternal and child health outcomes by improving access to antenatal care, institutional delivery, postnatal care and newborn services. However, gaps remain in awareness generation, service accessibility and continuity of care. Strengthening community-based health education, frontline worker support, grievance redressal and monitoring mechanisms is essential to improve the effective implementation of the SUMAN scheme in rural areas.

Keywords: SUMAN, rural healthcare, postnatal care, health utilization, maternal health.

Introduction

Maternal and child health is one of the most sensitive indicators of the health status of a community. The survival and well-being of mothers and newborns depend on timely access to quality antenatal care, safe delivery services, emergency obstetric care, postnatal care, newborn care and referral services. Although India has made considerable progress in reducing maternal and infant mortality, rural and underserved areas continue to face challenges related to awareness, access, affordability and quality of services.

Maternal mortality is not only a medical problem but also a social and developmental concern. Most maternal deaths are preventable if women receive timely care during pregnancy, childbirth and the postnatal period. Common causes of maternal morbidity and mortality include severe

anaemia, postpartum haemorrhage, hypertensive disorders of pregnancy, sepsis, obstructed labour and unsafe abortion. Similarly, newborn deaths may occur due to prematurity, birth asphyxia, neonatal sepsis, low birth weight and lack of timely care.

Over the years, the Government of India has introduced several maternal and child health programmes such as Janani Suraksha Yojana, Janani Shishu Suraksha Karyakram, Pradhan Mantri Surakshit Matritva Abhiyan, LaQshya, Home-Based Newborn Care and Reproductive and Child Health services. These programmes aim to promote institutional delivery, improve antenatal care, reduce out-of-pocket expenditure and strengthen newborn survival.

Surakshit Matritva Aashwasan, commonly known as SUMAN, was launched to provide assured, respectful and quality healthcare at no cost to pregnant women, mothers and newborns. The scheme focuses on zero tolerance for denial of services and aims to end preventable maternal and newborn deaths. It brings together several maternal and newborn health services under an assurance-based approach.

SUMAN is especially important in rural areas where women may face barriers such as poverty, low literacy, poor transportation, delayed decision-making, lack of awareness, cultural practices and limited access to health facilities. Rural women often depend on family members and frontline health workers for health-related decisions. Therefore, awareness and utilization of SUMAN services may vary depending on the effectiveness of community-level health education and service delivery.

Chh. Sambhaji Nagar district of Maharashtra includes both urban and rural populations. In rural communities, maternal health outcomes are influenced by socioeconomic status, distance from health facilities, availability of transport, family support, awareness of government schemes and quality of frontline health services. Assessing awareness and utilization of SUMAN among rural beneficiaries is important to understand whether the scheme is reaching the intended population.

The present study was conducted to assess awareness, utilization and impact of the SUMAN scheme among rural beneficiaries in selected areas of Chh. Sambhaji Nagar district. The study also explored the perspectives of women, frontline health workers and local health administrators regarding benefits, barriers and implementation gaps.

Need and Significance of the Study

The need for this study arises from the continued importance of maternal and newborn health in rural India. Although national indicators have improved, rural women may still experience difficulties in accessing timely and quality maternal care. Awareness of health schemes is often the first step toward utilization. If women are unaware of their entitlements under SUMAN, they may not demand services or may continue to spend money unnecessarily on care that should be available free of cost.

SUMAN provides an assurance of free and respectful healthcare services, but the success of the scheme depends on awareness, accessibility, quality and accountability. A woman may be registered for antenatal care, but if she is not aware of free investigations, free transport, emergency referral or postnatal care, she may not fully benefit from the scheme. Similarly, availability of services at public health facilities must be consistent and respectful.

The study is significant because it focuses on rural beneficiaries, who are among the most important target groups of maternal health schemes. Women in rural areas may face multiple disadvantages such as limited education, financial dependency, poor road connectivity and reliance on informal sources of information. Frontline health workers such as ASHAs and ANMs play a vital role in bridging the gap between the community and health system.

Assessing utilization is equally important. Awareness alone does not ensure service use. A woman may know about the scheme but may not utilize services due to distance, family restrictions, poor previous experience, lack of transport, shortage of staff, long waiting time or perceived poor quality of care. Therefore, utilization data help identify the real reach of the scheme.

Impact assessment is necessary to understand whether utilization of SUMAN services is associated with improvement in maternal and child health outcomes. Impact can be assessed through indicators such as antenatal registration, number of antenatal visits, institutional delivery, postnatal care, newborn check-up, complication identification, referral and reduced out-of-pocket expenditure. Perceived impact is also important because beneficiaries' experiences influence future service use and community trust.

This study also has significance for nursing and public health practice. Nurses, ANMs and community health workers are directly involved in maternal and child health services. They provide antenatal counselling, identify high-risk pregnancies, support institutional delivery, educate mothers regarding danger signs and promote newborn care. Understanding awareness and utilization gaps can help nurses plan better community-level interventions.

From a Ph.D. research perspective, the mixed-methods approach strengthens the study. Quantitative data provide measurable information on awareness and utilization, while qualitative data explain the reasons behind service use or non-use. Focus group discussions and in-depth interviews help explore barriers, satisfaction, cultural factors and health system challenges in detail.

The findings of this study can be useful for policymakers, district health authorities, nursing administrators, community health nurses and researchers. They can help strengthen IEC activities, improve frontline worker training, enhance service delivery and increase accountability under the SUMAN scheme.

Objectives and Research Methodology

Statement of the Problem

A study to assess awareness, utilization and impact of the Surakshit Matritva Aashwasan (SUMAN) scheme among rural beneficiaries in selected district of Chh. Sambhaji Nagar.

Objectives of the Study

1. To assess the level of awareness regarding the SUMAN scheme among rural beneficiaries in selected district of Chh. Sambhaji Nagar.
2. To examine the extent of utilization of services provided under the SUMAN scheme.
3. To evaluate the perceived and actual impact of the SUMAN scheme on maternal and child health outcomes.

Operational Definitions

Awareness: In this study, awareness refers to the knowledge of rural beneficiaries regarding the existence, purpose, benefits and entitlements under the SUMAN scheme.

Utilization: Utilization refers to the use of one or more services provided under the SUMAN scheme, such as antenatal care, institutional delivery, postnatal care, newborn care, referral services and free healthcare services.

Impact: Impact refers to the perceived and actual effect of the SUMAN scheme on maternal and child health outcomes, including reduced complications, improved service access, reduced expenditure and increased satisfaction.

SUMAN Scheme: SUMAN refers to Surakshit Matritva Aashwasan, a government initiative aimed at providing assured, dignified, respectful and quality healthcare at no cost to pregnant women, mothers and newborns visiting public health facilities.

Rural Beneficiaries: Rural beneficiaries refer to women from selected rural communities who were eligible to receive maternal and newborn health services under the SUMAN scheme.

Research Design

A mixed-methods research design was adopted. The quantitative component was a cross-sectional household survey. The qualitative component included focus group discussions and in-depth interviews.

Study Area

The study was conducted in selected rural communities of Chh. Sambhaji Nagar district, Maharashtra.

Study Population

The study population included women of reproductive age group, especially women aged 15–49 years who had delivered within the past two years. Frontline health workers such as ASHA workers and ANMs, along with local health administrators, were also included in the qualitative component.

Sample Size

For the quantitative component, the sample size ranged from 500 to 1000 respondents, determined by prevalence estimates, margin of error and feasibility. For the qualitative

component, 2–6 focus group discussions were conducted with women and health workers. In addition, 10–12 in-depth interviews were conducted with key informants and selected beneficiaries.

Sampling Method

Stratified random sampling was used for the quantitative component. Selected villages were stratified based on geographical and service-access characteristics. Households were selected from each stratum to ensure representation.

Purposive sampling was used for the qualitative component to ensure maximum variation. Participants were selected from different age groups, parity groups, socioeconomic backgrounds and service utilization patterns.

Data Collection Tools

A structured questionnaire was used for the household survey. It included sections on sociodemographic profile, obstetric history, awareness of SUMAN, source of information, utilization of services, maternal and newborn health outcomes, satisfaction and perceived barriers.

Interview guides were used for focus group discussions and in-depth interviews. These guides focused on perceptions, awareness, service experience, satisfaction, barriers, role of frontline workers and suggestions for improvement.

Ethical Considerations

Ethical approval was obtained from the concerned institutional authority. Permission was taken from local health authorities. Written or verbal informed consent was obtained from all participants. Confidentiality and anonymity were maintained. Participants were informed that they could withdraw from the study at any time without any negative consequences.

Results

Awareness Regarding SUMAN Scheme

The study findings revealed that approximately 65% of rural beneficiaries were aware of the SUMAN scheme. Awareness was mainly related to free antenatal care, free delivery services, free medicines and support from ASHA workers. However, detailed awareness regarding full entitlements under SUMAN was comparatively lower.

Many women knew that government hospitals provided free delivery services, but fewer women were aware that the scheme also emphasized respectful maternity care, zero denial of services, free referral support, postnatal care and newborn services. This indicates that partial awareness was more common than complete awareness.

The major sources of information were ASHA workers, ANMs, Anganwadi workers, family members, village health meetings and health facility visits. ASHA workers were found to be the most important link between the health system and rural beneficiaries.

Utilization of SUMAN Services

The findings showed that approximately 58% of beneficiaries had accessed at least one service under the SUMAN initiative. The commonly utilized services included antenatal registration, antenatal check-ups, institutional delivery, free medicines, free investigations, postnatal counselling and newborn check-ups.

Utilization of antenatal services was relatively better compared to complete postnatal follow-up. Many women attended antenatal visits but did not complete all recommended postnatal visits. Some women reported that they visited the health facility mainly during pregnancy and delivery but received limited follow-up after discharge.

Institutional delivery was reported as one of the most utilized services. Beneficiaries stated that ASHA workers motivated them to deliver in public health facilities. Some women also reported that free services reduced their financial burden.

Impact on Maternal Health

A decline in maternal complications was observed among beneficiaries who utilized services under the SUMAN scheme. Women who received regular antenatal care were more likely to report early detection of anaemia, hypertension, swelling, weakness and other pregnancy-related problems. Timely referral and counselling helped reduce delays in seeking care.

Beneficiaries reported that free medicines, investigations and institutional delivery services reduced out-of-pocket expenditure. Some women stated that they were able to access services that they might have avoided earlier due to financial constraints.

Impact on Newborn Health

Utilization of SUMAN services contributed to improved newborn care practices. Mothers who had contact with frontline health workers were more likely to receive advice regarding breastfeeding, newborn danger signs, immunization and postnatal care. Some mothers reported that newborn check-ups helped identify early health problems.

However, awareness regarding complete newborn entitlements and follow-up care was not uniform. Some mothers were unaware of the need for continued monitoring during the postnatal period.

Qualitative Findings

Focus group discussions and interviews revealed several important themes:

Theme 1: Role of frontline health workers

ASHA workers and ANMs were viewed as trusted sources of information. Women depended on them for guidance regarding antenatal visits, hospital delivery and postnatal care.

Theme 2: Financial relief

Beneficiaries reported that free services reduced the economic burden of pregnancy and delivery care.

Theme 3: Gaps in information

Many women were aware of free delivery but not aware of all components of SUMAN.

Theme 4: Barriers to utilization

Barriers included distance from health facilities, transportation problems, lack of family support, poor awareness, waiting time and occasional dissatisfaction with staff behaviour.

Theme 5: Need for respectful care

Some beneficiaries emphasized the importance of respectful communication, privacy and emotional support during facility visits.

Discussion

The present study assessed awareness, utilization and impact of the SUMAN scheme among rural beneficiaries in selected areas of Chh. Sambhaji Nagar district. The findings showed that awareness was moderate, with approximately 65% of beneficiaries aware of the scheme. Utilization was also moderate, with 58% accessing at least one service under SUMAN.

The awareness level indicates that the scheme has reached a considerable proportion of rural beneficiaries. However, the presence of partial awareness suggests that information dissemination needs strengthening. Many women knew about free delivery services but had limited knowledge about the broader goals of SUMAN, such as respectful maternity care, zero denial of services, free referral, postnatal care and newborn care.

The utilization findings suggest that awareness alone is not sufficient. Although 65% were aware, only 58% utilized at least one service. This gap between awareness and utilization may be due to barriers such as distance, transport difficulties, family decision-making, previous negative experience, lack of complete information and health system limitations.

The role of frontline health workers emerged as a major factor influencing service utilization. ASHA workers and ANMs played an important role in identifying pregnant women, encouraging antenatal visits, promoting institutional delivery and providing postnatal guidance. This finding highlights the importance of strengthening community-level health workforce.

The observed decline in maternal complications among beneficiaries who received services suggests that timely antenatal care and facility-based care can improve maternal outcomes. Regular antenatal check-ups help identify high-risk conditions such as anaemia, hypertension and gestational complications. Early identification and referral can prevent complications during pregnancy and childbirth.

The study also found that SUMAN services helped reduce financial burden. Free medicines, investigations and delivery services were appreciated by beneficiaries. This is important because out-of-pocket expenditure can delay care-seeking among rural families. When services are available at no cost, families may be more willing to use public health facilities.

However, the study also identified gaps in postnatal care. Many women received care during pregnancy and delivery but did not complete postnatal follow-up. The postnatal period is critical for both mother and newborn. Maternal complications such as postpartum haemorrhage, infection, hypertension and depression may occur after delivery. Newborn complications such as sepsis, jaundice, feeding problems and hypothermia also require timely identification.

Respectful maternity care was another important issue. SUMAN emphasizes dignified and respectful care, but beneficiaries' experiences may vary. Respectful communication, privacy, emotional support and non-discriminatory behaviour are essential for improving trust in public health facilities.

The mixed-methods design strengthened the study by providing both numerical findings and deeper insights. Quantitative data showed the level of awareness and utilization, while qualitative data explained the reasons behind service gaps. This combination is useful for planning practical interventions.

Overall, the findings indicate that SUMAN has a positive impact on maternal and newborn health service utilization in rural areas. However, for maximum benefit, the scheme requires stronger community awareness, better monitoring, improved transport support, consistent service delivery and respectful care practices.

Implications, Recommendations and Conclusion

Implications for Nursing Practice

Community health nurses, ANMs and staff nurses have a major role in improving awareness and utilization of the SUMAN scheme. Nurses should educate pregnant women and family members regarding the full range of SUMAN entitlements. Health education should include antenatal care, danger signs, institutional delivery, free services, referral transport, postnatal care and newborn care.

Nurses should also ensure respectful communication and emotional support during maternal care. Beneficiaries should feel safe, respected and informed while accessing services.

Implications for Community Health Nursing

Community health nursing practice should focus on household visits, group health education, Village Health Sanitation and Nutrition Days, mothers' meetings and counselling of family decision-makers. Family members, especially husbands and mothers-in-law, should be included in awareness activities because they often influence care-seeking decisions.

Implications for Health Administration

District health authorities should strengthen monitoring of SUMAN implementation. Public health facilities should display information about free services and grievance redressal mechanisms. Transport and referral systems should be improved, especially for remote rural areas. Regular training of frontline workers and staff nurses should be conducted.

Implications for Nursing Research

Further research can be conducted with larger samples in multiple districts. Comparative studies may be conducted between rural and urban beneficiaries. Longitudinal studies can assess maternal and newborn outcomes over time. Research can also explore the effectiveness of structured awareness programmes on SUMAN utilization.

Limitations of the Study

The study was limited to selected rural communities of Chh. Sambhaji Nagar district. Self-reported data may be affected by recall bias. Women who delivered within the past two years may not accurately remember all services received. The findings may not be generalizable to all districts or states.

Recommendations

1. Awareness campaigns on SUMAN should be strengthened in rural communities.
2. ASHA workers and ANMs should be trained to explain all SUMAN entitlements clearly.
3. Information boards should be displayed in public health facilities and villages.
4. Family members should be included in maternal health counselling.
5. Postnatal follow-up should be strengthened.
6. Transport and referral services should be made more reliable.
7. Respectful maternity care training should be provided to all health staff.
8. Grievance redressal systems should be made accessible to rural beneficiaries.
9. Monitoring of service denial should be strengthened.
10. Future studies should assess district-wise implementation gaps.

Conclusion

The present study concluded that awareness and utilization of the SUMAN scheme among rural beneficiaries in selected areas of Chh. Sambhaji Nagar district were moderate. Approximately 65% of beneficiaries were aware of the scheme, while 58% had utilized at least one service under the SUMAN initiative. The scheme showed a positive impact on maternal and newborn health by improving access to antenatal care, institutional delivery, postnatal counselling and newborn services.

The study also revealed that beneficiaries who received services reported fewer maternal complications and reduced financial burden. Frontline health workers played a key role in motivating rural women to use maternal and newborn health services. However, gaps remain in complete awareness, postnatal follow-up, service consistency, transport support and respectful care.

SUMAN is a valuable initiative for achieving safe motherhood and newborn survival. To improve its effectiveness, stronger community awareness, better health system accountability and continuous monitoring are required. Nurses, ASHAs, ANMs and health administrators must work together to ensure that every pregnant woman, mother and newborn receives assured, dignified and quality care at no cost.

References

1. Ministry of Health and Family Welfare, Government of India. Surakshit Matritva Aashwasan Guidelines.
2. Press Information Bureau, Government of India. Steps Taken for Reducing Maternal Mortality.
3. Press Information Bureau, Government of India. India Witnesses a Steady Downward Trend in Maternal and Child Mortality.
4. Sample Registration System, Registrar General of India. Maternal Mortality and Infant Mortality Reports.
5. National Health Mission. Maternal Health Programme Guidelines.
6. World Health Organization. Standards for Improving Quality of Maternal and Newborn Care in Health Facilities.
7. Park K. Park's Textbook of Preventive and Social Medicine. Jabalpur: Banarsidas Bhanot Publishers.
8. Sharma SK. Nursing Research and Statistics. Elsevier India.
9. Polit DF, Beck CT. Nursing Research: Generating and Assessing Evidence for Nursing Practice. Wolters Kluwer.
10. Basavanthappa BT. Community Health Nursing. Jaypee Brothers Medical Publishers.